## **Open Access**



Correction: Follicular reconstruction and neo-oogenesis in xenotransplantation of human ovarian isolated cells derived from chemotherapy-induced POF patients

Sara Khaleghi<sup>1</sup>, Farideh Eivazkhani<sup>2</sup>, Somayeh Tavana<sup>2</sup>, Ashraf Moini<sup>3,4</sup>, Marefat Ghaffari Novin<sup>1</sup>, Petkov Stoyan<sup>5,6</sup>, Hamid Nazarian<sup>1\*</sup> and Rouhollah Fathi<sup>2\*</sup>

## Correction: Journal of Biological Engineering (2023) 17:70

https://doi.org/10.1186/s13036-023-00384-2

The original article [1] unfortunately contained an error in the caption of Fig. 4: "IHC" should be corrected to "ICC", in order for the caption to read:

Fig. 4 ICC analysis of the HOCCs isolated from transsexual (A) and Chemo-POF (B) ovary. Anti-human Vimentin ICC was used to localize and quantify human stromal cells in the HOCCs, Anti-human Inhibin  $\alpha$  ICC was

The online version of the original article can be found at https://doi. org/10.1186/s13036-023-00384-2.

\*Correspondence: Hamid Nazarian H.Nazarian@sbmu.ac.ir Rouhollah Fathi rfathi79@royaninstitute.org <sup>1</sup>Department of Biology and Anatomical Sciences, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran <sup>2</sup>Department of Embryology, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran <sup>3</sup>Department of Endocrinology and Female Infertility, Royan Institute of Reproductive Biomedicine, ACECR, Tehran, Iran <sup>4</sup>Breast Disease Research Center (BDRC), Tehran University of Medical Science, Tehran, Iran <sup>5</sup>Platform Degenerative Diseases, German Primate Center, GmbH, Leibniz

Institute for Primate Research, 37077 Göttingen, Germany <sup>6</sup>German Center for Cardiovascular Research (DZHK), Partner Site

Göttingen, 37077 Göttingen, Germany

applied to localize and quantify human granulosa cells in the HOCCs and Anti-human Fragilis ICC was used to localize and quantify human OSCs in the HOCCs. C ICC analysis of OSCs isolated from transsexual ovary, after two passages, OSCs had increased in number. D Cell component comparison between Trans and Chemo-POF groups in passage 1 and passage 2. There are no significant differences in Vimentin positive cells between the groups in both passages 1 and 2. \*: The Inhibin  $\alpha$  and Fragilis positive cells have been significantly increased after passage 2 in both Trans and Chemo-POF groups (T1 with T2, T1 with P1, T2 with P2 and P1 with P2). E Alkaline phosphatase staining of OSCs. Scale bars: 50 µm. T1: Trans Passage 1; T2: Trans Passage 2, P1: POF Passage 1, P2: POF Passage 2.

The original article has been corrected. Published online: 07 October 2024

## Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/jublicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.